

**INFORMATION DISCLOSURE
CITATION**

(Use several sheets if necessary)

Atty. Docket No.

Serial No.

2425-18

09/973,568

Applicant

MANTZARIDIS et al

Filing Date

TC/A.U.

October 9, 2001

3736


U.S. PATENT DOCUMENTS

| *EXAMINER INITIAL | DOCUMENT NUMBER | DATE | NAME | CLASS | SUBCLASS | FILING DATE IF APPROPRIATE |
|----------------------|-----------------|------|------|-------|----------|-------------------------------|
| | | | | | | |
| | | | | | | |
| | | | | | | |

FOREIGN PATENT DOCUMENTS

| | | | | | | | TRANSLATION | |
|----------|--|--|------|---------|-------|----------|-------------|----|
| DOCUMENT | | | DATE | COUNTRY | CLASS | SUBCLASS | YES | NO |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |

OTHER DOCUMENTS (including Author, Title, Date, Pertinent pages, etc.)

| | | |
|----|----|---|
| JF | 1 | SEBEL et al., "The Cebrel Function Analysis Monitor (CFAM): A New Microprocessor-based Device for the On-line Analysis of the EEG and Evoked Potentials," British Journal of Anaesthesia (1983), Vol. 55, No. 12, pp. 1265-1270 |
| | 2 | KOCHS, E., "Electrophysiological monitoring and mild hypothermia," Journal of Neurosurgical Anaesthesiology (1995), Vol. 7, No. 3, pp. 222-228 |
| | 3 | SCHWENDER et al., "Spectral edge frequency of the electroencephalogram to monitor "depth" of anaesthesia with isoflurane or propofol," British Journal of Anaesthesia (1996), Vol. 77, No. 2, pp. 179-184 |
| | 4 | LESLIE et al., "Prediction of movement during propofol/nitrous oxide anesthesia," Anesthesiology (1996), Vol. 84, No. 1; pp. 52-63 |
| | 5 | GAITINI et al., "Awareness detection during caesarean section under general anaesthesia using EEG spectrum analysis," Canadian Journal of Anaesthesia (1995), Vol. 42, No. 5, pp. 37-381 |
| | 6 | TRAAST et al., "Electroencephalographic characteristics of emergence from propofol/sufentanil total intravenous anesthesia," Anesthesia and Analgesia (1995), Vol. 81, No. 2, pp. 366-371 |
| | 7 | SCHWILDEN et al., "Closed-loop feedback control of propofol anaesthesia by quantitative EEG analysis in humans," British Journal of Anaesthesiology," (1989), Vol. 62, No. 3, pp. 290-296 |
| | 8 | RUSS et al., "Spectral analysis of the EEG during hypothermic cardiopulmonary bypass," Acta Anaesthesiologica Scandinavica (1987), Vol. 31, pp. 111-116 |
| | 9 | LEVY, W.J., "Quantitative analysis of EEG changes during hypothermia," Anesthesiology (1984), Vol. 60, No. 4, pp. 291-297 |
| | 10 | BASHEIN et al., "Electroencephalography during surgery with cardiopulmonary bypass and hypothermia," Anesthesiology (1992), Vol. 76, No. 6, pp. 878-891 |
| | 11 | FLAISHON et al., "Detection of consciousness following thiopental: isolated forearm and bispectral EEG (BIS)," Anesthesiology (1995), Vol. 83, No. 3A, A515 |
| | 12 | KEARSE et al., "The Bispectral Index correlates with sedation/hypnosis and recall: comparison using multiple agents," Anesthesiology (1995), Vol. 83, No. 3A, A507 |
| | 13 | KEARSE et al., "Bispectral analysis of the electroencephalogram correlates with patient movement to skin incision during propofol/nitrous oxide anesthesia," Anesthesiology (1994), Vol. 81, No. 6, pp. 1365-1370 |
| JF | 14 | VERNON et al., "Prediction of movement using bispectral electroencephalographic analysis during propofol/alfentanil or isoflurane/alfentanil anesthesia," Anesthesia & Analgesia (1995), Vol. 80, No. 4, pp. 780-785 |

*Examiner

Date Considered

10/15/04

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to application.

INFORMATION DISCLOSURE
CITATION

(Use several sheets if necessary)

Atty. Docket No.

2425-18

Applicant

MANTZARIDIS et al

Filing Date

October 9, 2001

Serial No.

09/973,568

TC/A.U.

3736



OTHER DOCUMENTS (including Author, Title, Date, Pertinent pages, etc.)

| | |
|----|---|
| 15 | THORNTON et al., "Effect of propofol on the auditory evoked response and oesophageal contractility," British Journal of Anaesthesia (1989), Vol. 63, No. 4, pp. 411-417 |
| 16 | DAVIES et al., "Middle latency auditory evoked potentials during repeated transitions from consciousness to unconsciousness," Anaesthesia (1996), Vol. 51, No. 1, pp. 107-113 |
| 17 | NEWTON et al., "Auditory evoked response and awareness: a study of volunteers at sub-MAC concentrations of isoflurane," British Journal of Anaesthesia (1992), Vol. 69, No. 2, pp. 122-129 |
| 18 | KENNY et al., "Transition between consciousness and unconsciousness during anesthesia," Anesthesiology (1993), Vol. 79, No. 3A, A330 |
| 29 | KENNY et al., "Closed-loop control of anesthesia," Anesthesiology (1992), Vol. 77, No. 3A, A328 |
| 20 | HETT et al., "Effect of temperature and cardiopulmonary bypass on the auditory evoked response," British Journal of Anaesthesia (1995), Vol. 75, pp. 293-296 |
| 21 | KENNY et al., "A portable target controlled propofol infusion system," International Journal of Clinical Monitoring and Computing (1992), Vol. 9, No. 3, pp. 179-182 |
| 22 | DAVIES et al., "Postoperative analgesia using a computerized infusion of alfentanil following aortic bifurcation graft surgery," International Journal of Clinical Monitoring and Computing (1992), Vol. 9, No. 4, pp. 207-212 |
| 23 | MARKAND et al., "Monitoring of multimodality evoked potentials during open heart surgery under hypothermia," Electroencephalography and Clinical Neurophysiology (1984), Vol. 59, No. 6, pp. 432-440 |
| 24 | BLAIR, E., "A physiological classification of clinical hypothermia," Surgery (1965), Vol. 58, No. 3, pp. 607-618 |
| 25 | CHASSARD et al., "Auditory evoked potentials during propofol anaesthesia in man," British Journal of Anaesthesia (1989), Vol. 62, No. 5, pp. 522-526 |
| 26 | LESLIE et al., "Propofol blood concentration and the Bispectral Index predict suppression of learning during propofol/epidural anesthesia in volunteers," Anesthesia & Analgesia (1995) Vol. 81, No. 6, pp. 1269-1274 |
| 27 | DAVIDSON et al., "Effective concentration 50 for propofol with and without 67% nitrous oxide," Acta Anaesthesiologica Scandinavica (1993), Vol. 37, pp. 458-464 |
| 28 | RUSSELL et al., "Propofol-fentanyl anaesthesia for coronary artery surgery and cardiopulmonary bypass," Anaesthesia (1989), Vol. 44, No. 3, pp. 205-208 |
| 29 | MASSEY et al., "Pharmacokinetics of an infusion of propofol during cardiac surgery," British Journal of Anaesthesia (1990), Vol. 65, No. 4, pp. 475-479 |
| 30 | HYNEN et al., "Propofol sequestration within the extracorporeal circuit," Canadian Journal of Anaesthesia (1994), Vol. 41, No. 7, pp. 583-588 |
| 31 | SCHWENDER et al., "Effects of increasing doses of alfentanil, fentanyl and morphine on mid-latency auditory evoked potentials," British Journal of Anaesthesia (1993), Vol. 71, No. 5, pp. 622-628 |
| 32 | JESSOP et al., "Evaluation of the actions of general anaesthetics in the human brain," General Pharmacology (1992), Vol. 23, No. 6, pp. 927-935 |
| 33 | JONES, J.G., "Perception and memory during general anaesthesia," British Journal of Anaesthesia (1994), Vol. 73, No. 1, pp. 31-37 |
| 34 | SHARPE et al., "Auditory evoked response, median frequency and 95% spectral edge during anaesthesia with desflurane and nitrous oxide," British Journal of Anaesthesia (1997), Vol. 78, pp. 282-285 |
| 35 | KEARSE et al., "Bispectral analysis of the electroencephalogram during induction of anesthesia may predict hemodynamic responses to laryngoscopy and intubation," Electroencephalography & Clinical Neurophysiology (1994), Vol. 90, No. 3, pp. 194-200 |
| 36 | SEBEL et al., "EEG bispectrum predicts movement during thiopental/isoflurane anesthesia," Journal of Clinical Monitoring (1995), Vol. 11, No. 2, pp. 83-91 |
| 37 | LIU et al., "Electroencephalogram bispectral analysis predicts the depth of midazolam-induced sedation," Anesthesiology (1996), Vol. 84, No. 1, pp. 64-69 |

*Examiner

Date Considered

10/15/04

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to application.

**INFORMATION DISCLOSURE
CITATION**

(Use several sheets if necessary)

Atty. Docket No.

2425-18

Applicant

MANTZARIDIS et al


Filing Date

October 9, 2001

Serial No.

09/973,568

TC/A.U.

3736

| | |
|----|--|
| 38 | KENNY et al., "Validation of monitoring anesthetic depth by closed-loop control. In: Memory and Awareness in Anesthesia," Sebel P. Bonke, Winograd E. eds. Prentice Hall, Englewood Cliffs (1993), pp. 255-264 |
| 39 | LEVY et al., "Automated EEG processing for intraoperative monitoring: a comparison of techniques," Anesthesiology (1980), Vol. 53, No. 3, pp. 223-236 |
| 40 | SIGL et al., "An introduction to bispectral analysis for the electroencephalogram," Journal of Clinical Monitoring (1994), Vol. 10, No. 6, pp. 392-404 |
| 41 | HOWELL et al., "Defining the CP ₅₀ and BIS ₅₀ for propofol alone and propofol with alfentanil," Anesthesiology (1995), Vol. 83, No. 1, A367 |
| 42 | RUSSELL, I.F., "Midazolam-alfentanil: an anaesthetic? An investigation using the isolated forearm technique," British Journal of Anaesthesia (1993), Vol. 70, No. 1, pp. 42-46 |
| 43 | LAMBRECHTS et al., "Postoperative amnesia," British Journal of Anaesthesia (1961), Vol. 33, No. 8, pp. 397-404 |
| 44 | ARTUSIO, J.F. Jr., "Ether analgesia during major surgery," Journal of the American Medical Association (1955), Vol. 157, No. 1, pp. 33-36 |
| 45 | RUPREHT, J., "Awareness with amnesia during total intravenous anaesthesia with propofol (letter)," Anaesthesia (1989), Vol. 44, No. 12, p. 1005 |
| 46 | SAWTELLE et al., "Bispectral EEG index predicts awakening," Anesthesiology (1994), Vol. 81, No. 3A, A213 |
| 47 | GAJRAJ et al., "A comparison of auditory evoked potentials and bispectral EEG analysis in spontaneously breathing anesthetized patients," Anesthesiology (1996), Vol. 85, No. 3A, A462 |
| 48 | SCHWILDEN et al., "Quantitative EEG analysis during anaesthesia with isoflurane in nitrous oxide at 1.3 and 1.5 MAC," British Journal of Anaesthesia (1987), Vol. 59, No. 6, pp. 738-745 |
| 49 | SCHWILDEN et al., "Closed-loop feedback control of methohexital anesthesia by quantitative EEG analysis in humans," Anesthesiology (1987), Vol. 67, No. 3, pp. 341-347 |
| 50 | SCHWILDEN et al., "Quantitation of the EEG and pharmacodynamic modeling of hypnotic drugs: etomidate as an example," European Journal of Anaesthesiology (1995), Vol. 2, No. 2, pp. 121-131 |
| 51 | LEVY, W.I., "Intraoperative EEG patterns: implications for EEG monitoring," Anesthesiology (1984), Vol. 60, No. 5, pp. 430-434 |
| 52 | LIU et al., "Incidence of awareness with recall during general anesthesia," Anaesthesia (1991), Vol. 46, No. 6, pp. 435-437 |
| 53 | ARNDT et al., "EEG-Veränderungen unter Propofol-Alfentanil-Lachgas-Naekoaw," [EEG changes during propofol-alfentanil-nitrous oxide anesthesia] [German] (1995), Vol. 20, pp. 126-133 |
| 54 | CLARK et al., "Neurophysiologic effects of general anesthetics: I. The electroencephalogram and sensory evoked responses in man," Anesthesiology (1973), Vol. 38, No. 6, pp. 564-582 |
| 55 | ROSNER et al., "Neurophysiologic effects of general anesthetics: II Sequential regional actions in the brain," Anesthesiology (1973), Vol. 39, No. 1, pp. 59-81 |
| 56 | VERNON et al., "EEG bispectrum predicts movement at incision during isoflurane or propofol anesthesia," Anesthesiology (1992), Vol. 77, No. 3A, A502 |
| 57 | GLASS et al., "Quantification of the relative effects of anesthetics agents on the EEG and patient responsiveness to incision," Anesthesiology (1994), Vol. 81, No. 3A, A407 |
| 58 | LANG et al., "Bispectral EEG analysis, analgesia and movement at incision during propofol/alfentanil/N2O anesthesia," Anesthesiology (1994), Vol. 81, No. 3A, A476 |
| 59 | SEBEL et al., "Bispectral analysis (BIS) for monitoring anesthesia: comparison of anesthetic techniques," Anesthesiology (1994), Vol. 81, No. 3A, A1488 |
| 60 | THORNTON et al., "Effects of halothane or enflurane with controlled ventilation on auditory evoked potentials," British Journal of Anaesthesia (1984), Vol. 56, No. 4, pp. 315-323 |

*Examiner

Date Considered

10/15/04

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to application.

INFORMATION DISCLOSURE
CITATION

(Use several sheets if necessary)

Atty. Docket No.

2425-18

Applicant

MANTZARIDIS et al

Filing Date

October 9, 2001

Serial No.

09/973,568

TC/A.U.

3736



| OTHER DOCUMENTS (including Author, Title, Date, Pertinent pages, etc.) | |
|--|---|
| 61 | THORNTON et al., "Effect of etomidate on the auditory evoked response in man," British Journal of Anaesthesia (1985), Vol. 57, No. 6, pp. 554-561 |
| 62 | HENEGHAN et al., "Effect of isoflurane on the auditory evoked response in man," British Journal of Anaesthesia (1987), Vol. 59, No. 3, pp. 277-282 |
| 63 | de BEER et al., "Haemodynamic responses to incision and sternotomy in relation to the auditory evoked potential and spontaneous EEG," British Journal of Anaesthesia (1996), Vol. 76, No. 5, pp. 685-693 |
| 64 | THORNTON et al., "Effects of surgical stimulation on the auditory evoked response," British Journal of Anaesthesia (1988), Vol. 60, No. 4, pp. 372-378 |
| 65 | SCHWENDER et al., "Effects of surgical stimulation on midlatency auditory evoked potentials during general anaesthesia with propofol/fentanyl, isoflurane/fentanyl and flunitrazepam/fentanyl," Anaesthesia (1994), Vol. 49, No. 7, pp. 572-578 |
| 66 | SEBEL et al., "Bispectral analysis for monitoring anesthesia - a multicenter study," Anesthesiology (1993), Vol. 79, No. 3A, A178 |
| 67 | THORNTON et al., "Enflurane anaesthesia causes graded changes in the brainstem and early cortical auditory evoked response in man," British Journal of Anaesthesia (1983), Vol. 55, No. 6, pp. 479-486 |
| 68 | THORNTON et al., "Selective effect of althesin on the auditory evoked response in man," British Journal of Anaesthesia (1986), Vol. 58, No. 4, pp. 422-427 |
| 69 | GOLDSTEIN et al., "Effects of stimulus rate and number on the early components of the averaged electroencephalographic response," Journal of Speech and Hearing Research (1972), Vol. 15, No. 3, pp. 559-566 |
| 70 | McFARLAND et al., "Reexamination of effects of stimulus rate and number on the middle components of the averaged electroencephalographic response," Audiology (1975), Vol. 14, pp. 456-465 |
| 71 | THORNTON et al., "The auditory evoked response: a measure of depth of anaesthesia," Bailliere's Clinical Anesthesiology (1989), Vol. 3, No. 3, pp. 559-585 |
| 72 | SEBEL et al., "Evoked responses - a neurophysiological indicator of depth of anaesthesia? (editorial), British Journal of Anaesthesia (1985), Vol. 57, No. 9, pp. 841-842 |
| 73 | THORNTON, C., "Evoked potentials in anaesthesia," European Journal of Anaesthesiology (1991), Vol. 8, pp. 89-107 |
| 74 | MADLER et al., "Auditory evoked potentials indicate the loss of neuronal oscillations during general anaesthesia," Naturwissenschaften (1987), Vol. 74, pp. 42-43 |
| 75 | MADLER et al., "Sensory information processing during general anaesthesia: effect of isoflurane on auditory evoked neuronal oscillations," British Journal of Anaesthesia (1991), Vol. 66, No. 1, pp. 81-87 |
| 76 | SCHWENDER et al., "Mid-latency auditory evoked potentials during ketamine anaesthesia in humans," British Journal of Anaesthesia (1993), Vol. 71, No. 5, pp. 629-632 |
| 77 | SCHWENDER et al., "Motor signs of wakefulness during general anaesthesia with propofol, isoflurane and flunitrazepam/fentanyl and midlatency auditory evoked potentials," Anaesthesia (1994), Vol. 49, No. 6, pp. 476-484 |
| 78 | THORNTON et al., "The auditory evoked response as an indicator of awareness," British Journal of Anaesthesia (1989), Vol. 63, No. 1, pp. 113-115 |

*Examiner

Date Considered

10/15/04

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to application.